


# SIMPLICIAL SPACES



CS 468 – Lecture 3


8/10/2

# OVERVIEW

- First lecture: Point set topology
  - Manifolds: locally Euclidean
  - Homeomorphisms: bijective bi-continuous maps
  - Classification Theorem for 2-Manifolds
  - How to compute?
- This lecture: Combinatorial topology
  - Simplicial Complexes
  - Subcomplexes
  - Triangulations and Orientability
  - Euler Characteristic
  - 2-Manifold Homeomorphism problem (revisited)

## GEOMETRIC DEFINITION: SIMPLEX

- A ***k*-simplex** is the convex hull of  $k + 1$  affinely independent points  $S = \{v_0, v_1, \dots, v_k\}$ . The points in  $S$  are the **vertices** of the simplex.
- A  $k$ -simplex is a  $k$ -dimensional subspace of  $\mathbb{R}^d$ ,  $\dim \sigma = k$ .




## GEOMETRIC DEFINITION: FACES

- $\sigma$ : a  $k$ -simplex defined by  $S = \{v_0, v_1, \dots, v_k\}$ .
- $\tau$  defined by  $T \subseteq S$  is a **face** of  $\sigma$
- $\sigma$  is its **coface**.
- $\sigma \geq \tau$  and  $\tau \leq \sigma$ .
- $\sigma \leq \sigma$  and  $\sigma \geq \sigma$ .


vertex  
a

edge  
[a, b]

triangle  
[a, b, c]




tetrahedron  
[a, b, c, d]



## GEOMETRIC DEFINITION: SIMPLICIAL COMPLEX

- A **simplicial complex**  $K$  is a finite set of simplices such that
  1.  $\sigma \in K, \tau \leq \sigma \Rightarrow \tau \in K,$
  2.  $\sigma, \sigma' \in K \Rightarrow \sigma \cap \sigma' \leq \sigma, \sigma'$  or  $\sigma \cap \sigma' = \emptyset.$
- The **dimension** of  $K$  is  $\dim K = \max\{\dim \sigma \mid \sigma \in K\}.$
- The **vertices** of  $K$  are the zero-simplices in  $K$ .
- A simplex is **principal** if it has no proper coface in  $K$ .




## SIZE OF A SIMPLEX:


# LOW DIMENSIONS

vertex  
a

edge  
[a, b]



triangle  
[a, b, c]




tetrahedron  
[a, b, c, d]

$k/l$	0	1	2	3
0	1	0	0	0
1	2	1	0	0
2	3	3	1	0
3	4	6	4	1
4	?	?	?	?


## SIZE OF A SIMPLEX: ANOTHER VIEW

vertex  
a


edge  
[a, b]



triangle  
[a, b, c]





tetrahedron  
[a, b, c, d]



## SIZE OF A SIMPLEX:

# PASCAL'S TRIANGLE



## SIZE OF A SIMPLEX: BINOMIAL COEFFICIENTS

- $\emptyset$  is the **( $-1$ )-simplex**.
- A  $k$ -simplex has  $\binom{k+1}{l+1}$  faces of dimension  $l$
- Total size is:

$$\sum_{l=-1}^k \binom{k+1}{l+1} = 2^{k+1}$$

- Large, simplex, uniform

**ABSTRACT DEFINITION:**  
**SIMPLICIAL COMPLEX**

- An **abstract simplicial complex** is a set  $K$ , together with a collection  $\mathcal{S}$  of subsets of  $K$  called **(abstract) simplices** such that:
  1. For all  $v \in K$ ,  $\{v\} \in \mathcal{S}$ . We call the sets  $\{v\}$  the **vertices** of  $K$ .
  2. If  $\tau \subseteq \sigma \in \mathcal{S}$ , then  $\tau \in \mathcal{S}$ .
- We call  $\mathcal{S}$  the complex.




## ABSTRACT DEFINITION: RELATIONSHIP

- Let  $K$  be a simplicial complex with vertices  $V$  and let  $\mathcal{S}$  be the collection of all subsets  $\{v_0, v_1, \dots, v_k\}$  of  $V$  such that the vertices  $v_0, v_1, \dots, v_k$  span a simplex of  $K$ . Then,  $\mathcal{S}$  is the **vertex scheme** of  $K$ .
- $K$  and  $\mathcal{S}$  form an abstract simplicial complex.
- Two abstract simplicial complexes are **isomorphic** if we can one from the other by renaming vertices.
- (Theorem) Every abstract complex  $\mathcal{S}$  is isomorphic to the vertex scheme of some simplicial complex  $K$ .
- We call  $K$  a **geometric realization** of  $\mathcal{S}$ .

## WAVEFRONT's OBJ FORMAT

```
v -0.269616 0.228466 0.077226
v -0.358878 0.240631 0.044214
v -0.657287 0.527813 0.497524
v 0.186944 0.256855 0.318011
v -0.074047 0.212217 0.111664
...
f 19670 20463 20464
f 8936 8846 14300
f 4985 12950 15447
f 4985 15447 15448
...
```

## SUBCOMPLEXES: EXAMPLE




## SUBCOMPLEXES: POSETS

- Let  $S$  be a finite set. A **partial order** is a binary relation on  $S$  such that all  $x, y, z \in S$ ,
  1. (Reflexive)  $x \leq x$ ,
  2. (Antisymmetric)  $x \leq y$  and  $y \leq x$  implies  $x = y$ ,
  3. (Transitive)  $x \leq y$  and  $y \leq z$  implies  $x \leq z$ .
- A set with a partial order is a **partially ordered set** or **poset** for short.
- Partial vs. full orders


## SUBCOMPLEXES: CLOSURE

- A **subcomplex** is a simplicial complex  $L \subseteq K$ . The smallest subcomplex containing a subset  $L \subseteq K$  is its closure,  
 $\text{Cl } L = \{\tau \in K \mid \tau \leq \sigma \in L\}$ .
- Everything “below” is included.



## SUBCOMPLEXES: STAR


- The **star of  $L$**  contains all of the cofaces of  $L$ ,  
 $\text{St } L = \{\sigma \in K \mid \sigma \geq \tau \in L\}.$
- Everything “above” is included.
- Stars are analogs of neighborhoods (open).



## SUBCOMPLEXES: LINK


- The link of  $L$  is the boundary of its star,

$$\text{Lk } L = \text{Cl St } L - \text{St}(\text{Cl } L - \{\emptyset\}).$$



# TRIANGULATIONS

- The **underlying space**  $|K|$  of a simplicial complex  $K$  is  $|K| = \cup_{\sigma \in K} \sigma$ .
- $|K|$  is a topological space.
- A **triangulation** of a topological space  $\mathbb{X}$  is a simplicial complex  $K$  such that  $|K| \approx \mathbb{X}$ .







# ORIENTABILITY

- An **orientation** of a  $k$ -simplex  $\sigma \in K$ ,  $\sigma = \{v_0, v_1, \dots, v_k\}$ ,  $v_i \in K$  is an equivalence class of orderings of the vertices of  $\sigma$ , where

$$(v_0, v_1, \dots, v_k) \sim (v_{\tau(0)}, v_{\tau(1)}, \dots, v_{\tau(k)})$$

are equivalent orderings if the parity of the permutation  $\tau$  is even.

- We denote an **oriented simplex**, a simplex with an equivalence class of orderings, by  $[\sigma]$ .



# ORIENTABILITY

- Two  $k$ -simplices sharing a  $(k - 1)$ -face  $\sigma$  are **consistently oriented** if they induce different orientations on  $\sigma$ .
- A triangulable  $d$ -manifold is **orientable** if all  $d$ -simplices can be oriented consistently.
- Otherwise, the  $d$ -manifold is **non-orientable**

vertex  
a

edge  
[a, b]

triangle  
[a, b, c]

tetrahedron  
[a, b, c, d]

# INVARIANTS

- A **(topological) invariant** is a map  $f$  that assigns the same object to spaces of the same topological type.
- $\mathbb{X} \approx \mathbb{Y} \implies f(\mathbb{X}) = f(\mathbb{Y})$
- $f(\mathbb{X}) \neq f(\mathbb{Y}) \implies \mathbb{X} \not\approx \mathbb{Y}$  (contrapositive)
- $f(\mathbb{X}) = f(\mathbb{Y}) \implies$  nothing
- “coarser” differentiation


## EULER CHARACTERISTIC: DEFINITION

- $K$  a simplicial complex with  $s_k$   $k$ -simplices.
- The Euler characteristic  $\chi(K)$  is

$$\chi(K) = \sum_{i=0}^{\dim K} (-1)^i s_i = \sum_{\sigma \in K - \{\emptyset\}} (-1)^{\dim \sigma}.$$

- $v - e + f = 1$  (Graph Theory)
- Invariant for  $|K|$
- Any triangulation gives the same answer!
- Intrinsic property


## EULER CHARACTERISTIC: BASIC 2-MANIFOLDS



2-Manifold	$\chi$
Sphere $\mathbb{S}^2$	2
Torus $\mathbb{T}^2$	0
Klein bottle $\mathbb{K}^2$	0
Projective plane $\mathbb{RP}^2$	1

## EULER CHARACTERISTIC: CONNECTED SUMS

- (Theorem) For compact surfaces  $\mathbb{M}_1, \mathbb{M}_2$ ,  
$$\chi(\mathbb{M}_1 \# \mathbb{M}_2) = \chi(\mathbb{M}_1) + \chi(\mathbb{M}_2) - 2.$$
- $\chi(g\mathbb{T}^2) = 2 - 2g$
- $\chi(g\mathbb{R}\mathbf{P}^2) = 2 - g$
- The connected sum of  $g$  tori is called a surface with **genus**  $g$ .



## EULER CHARACTERISTIC: HOMEOMORPHISM PROBLEM

- (Theorem) Closed compact surfaces  $\mathbb{M}_1$  and  $\mathbb{M}_2$  are homeomorphic,  
 $\mathbb{M}_1 \approx \mathbb{M}_2$  iff
  1.  $\chi(\mathbb{M}_1) = \chi(\mathbb{M}_2)$  and
  2. either both surfaces are orientable or both are non-orientable.
- “iff” so full answer. We’re done!
- Higher dimensions?